

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> :

A61K 31/70, 35/80

A3

(11) International Publication Number:

WO 94/24984

(43) International Publication Date: 10 November 1994 (10.11.94)

(21) International Application Number: PCT/US94/04728

(22) International Filing Date: 29 April 1994 (29.04.94)

(30) Priority Data:

08/055,533

30 April 1993 (30.04.93)

US

(71)(72) Applicant and Inventor: WINGET, Rodney, R. [US/US];  
13265 - 89th Avenue South, Renton, WA 98055 (US).(74) Agents: HERMANN, Karl, R. et al.; Seed and Berry, 6300  
Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-  
7092 (US).(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FL,  
GE, HU, JP, KG, KP, KR, KZ, LK, LV, MD, MG, MN,  
MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, UZ,  
VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB,  
GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ,  
CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

## Published

*With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*


(88) Date of publication of the international search report:

2 February 1995 (02.02.95)

(54) Title: ANTI-INFLAMMATORY COMPOSITIONS CONTAINING EICOSAPENTAENOIC ACID BEARING MONOGALACTO-  
SYLDIACYLGLYCEROL

## (57) Abstract

There is disclosed an anti-inflammatory composition containing eicosapentaenoic acid (EPA) bearing monogalactosyldiacylglycerol (MGDG-EPA) at high levels. Additional components of the composition include digalactosyldiacylglycerol (DGDG), phosphatidylcholine (PC), chlorophylls and carotenoids. Methods for making the anti-inflammatory composition by extraction of marine algae, as well as methods of treating inflammation by administering an effective amount of the composition to an animal, are also disclosed.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |                                       |    |                          |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria                  | GB | United Kingdom                        | MR | Mauritania               |
| AU | Australia                | GE | Georgia                               | MW | Malawi                   |
| BB | Barbados                 | GN | Guinea                                | NE | Niger                    |
| BE | Belgium                  | GR | Greece                                | NL | Netherlands              |
| BF | Burkina Faso             | HU | Hungary                               | NO | Norway                   |
| BG | Bulgaria                 | IE | Ireland                               | NZ | New Zealand              |
| BJ | Benin                    | IT | Italy                                 | PL | Poland                   |
| BR | Brazil                   | JP | Japan                                 | PT | Portugal                 |
| BY | Belarus                  | KE | Kenya                                 | RO | Romania                  |
| CA | Canada                   | KG | Kyrgyzstan                            | RU | Russian Federation       |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan                    |
| CG | Congo                    | KR | Republic of Korea                     | SE | Sweden                   |
| CH | Switzerland              | KZ | Kazakhstan                            | SI | Slovenia                 |
| CI | Côte d'Ivoire            | LI | Liechtenstein                         | SK | Slovakia                 |
| CM | Cameroon                 | LK | Sri Lanka                             | SN | Senegal                  |
| CN | China                    | LU | Luxembourg                            | TD | Chad                     |
| CS | Czechoslovakia           | LV | Latvia                                | TG | Togo                     |
| CZ | Czech Republic           | MC | Monaco                                | TJ | Tajikistan               |
| DE | Germany                  | MD | Republic of Moldova                   | TT | Trinidad and Tobago      |
| DK | Denmark                  | MG | Madagascar                            | UA | Ukraine                  |
| ES | Spain                    | ML | Mali                                  | US | United States of America |
| FI | Finland                  | MN | Mongolia                              | UZ | Uzbekistan               |
| FR | France                   |    |                                       | VN | Viet Nam                 |
| GA | Gabon                    |    |                                       |    |                          |

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 94/04728

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 A61K31/70 A61K35/80

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No. |
|------------|--|-----------------------|
| Y          | US,A,5 077 202 (SETO ET AL.) 31 December 1991<br>see the whole document<br>---   | 1-12                  |
| Y          | DE,A,40 22 815 (B.BRAUN MELSUNGEN) 23 January 1992<br>see the whole document<br>---  | 1-12                  |
| Y          | CLINICAL RESEARCH,<br>vol.40, no.2, 1992<br>page 164A<br>G.G. KRUEGER ET AL. 'PERCUTANEOUS<br>ABSORPTION AND ANTI-INFLAMMATORY ACTIVITY<br>OF TOPICALLY APPLIED LIPIDS CONTAINING<br>EITHER ARACHIDONIC ACID OR<br>EICOSAPENTAENOIC ACID'<br>see the whole document<br>--- | 1-12                  |

--/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*a\* document member of the same patent family

Date of the actual completion of the international search

20 December 1994

Date of mailing of the international search report

06.01.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

## INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/US 94/04728

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| Y          | JOURNAL OF GENERAL MICROBIOLOGY,<br>vol.139, no.3, March 1993<br>pages 465 - 472<br>W. YONGMANITCHAI ET AL. 'POSITIONAL<br>DISTRIBUTION OF FATTY ACIDS, AND MOLECULAR<br>SPECIES OF POLAR LIPIDS, IN THE DIATOM<br>PHAEODACTYLUM TRICORNUTUM'<br>see the whole document<br>---  | 1-12                  |
| A          | BULLETIN OF THE JAPANESE SOCIETY OF<br>SCIENTIFIC FISHERIES,<br>vol.37, no.4, 1971<br>pages 326 - 332<br>S. SATO 'STUDIES ON GLYCOLIPIDS IN MARINE<br>ALGAE'<br>see the whole document<br>---   | 7-10                  |
| Y          | CHEMICAL & PHARMACEUTICAL BULLETIN,<br>vol.40, no.6, 1992<br>pages 1404 - 1410<br>M. KOBAYASHI ET AL. 'MARINE NATURAL<br>PRODUCTS. XXIX.'<br>see the whole document<br>---  | 1-12                  |
| Y          | CHEMICAL & PHARMACEUTICAL BULLETIN,<br>vol.37, no.3, 1989<br>pages 849 - 851<br>I. KITAGAWA ET AL.<br>'HETEROSIGMA-GLYCOLIPIDS I AND II, TWO NEW<br>GALACTOLIPIDS CONTAINING<br>OCTADECATETRAENOYL AND EICOSAPENTAENOYL<br>RESIDUES, FROM A RAPHIDOPHYTE<br>DINOFLAGELLATE HETEROSIGMA SP'<br>see the whole document<br>--- | 1-12                  |
| Y          | CHEMICAL & PHARMACEUTICAL BULLETIN,<br>vol.30, no.10, 1982<br>pages 3544 - 3547<br>H. KIRUCHI ET AL. 'MARINE NATURAL<br>PRODUCTS. X.'<br>see the whole document<br>---  | 1-12                  |
| Y          | DATABASE WPI<br>Week 8242,<br>Derwent Publications Ltd., London, GB;<br>AN 82-89599E<br>'GLYCOLIPID HAVING ANTIINFLAMMATORY AND<br>ANTIALLERGIC ACTIVITY'<br>& JP,A,57 149 299 (FUJISAWA PHARM) 14<br>September 1982<br>see abstract<br>---   | 1-12                  |
|            | ---   |                       |
|            | -/--  |                       |

## INTERNATIONAL SEARCH REPORT

Inter- national Application No  
PCT/US 94/04728

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No. |
|----------|--|-----------------------|
| Y        | PATENT ABSTRACTS OF JAPAN<br>vol. 15, no. 332 (C-0861) 23 August 1991<br>& JP,A,03 127 729 (KATO TAKAYOSHI) 30 May 1991<br>see abstract<br>---                                       | 1-12                  |
| A        | JAP. J. PHYCOL.,<br>vol.34, 1986<br>pages 94 - 100<br>S. ARAKI ET AL. 'LIPID AND FATTY ACID<br>COMPOSITION IN THE RED ALGA PORPHYRA<br>YEZOENSIS'<br>see the whole document<br>----- | 7-10                  |

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 94/ 04728

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11-12  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 11 and 12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/04728

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| US-A-5077202                              | 31-12-91            | JP-C- 1670333              | 12-06-92            |
|   |                     | JP-B- 3027538              | 16-04-91            |
|   |                     | JP-A- 61063624             | 01-04-86            |
| DE-A-4022815                              | 23-01-92            | DE-A- 4042437              | 11-06-92            |

### Description

#### ANTI-INFLAMMATORY COMPOSITIONS CONTAINING EICOSAPENTAENOIC ACID BEARING MONOGALACTOSYL- DIACYLGLYCEROL AND METHODS RELATING THERETO

### Technical Field

This invention relates generally to an anti-inflammatory composition and, more specifically, to an anti-inflammatory composition derived from marine algae having a high content of eicosapentaenoic acid bearing monogalactosyldiacylglycerol (MGDG-EPA).

### Background of the Invention


Monogalactosyldiacylglycerides (MGDG) may generally be obtained from a number of higher plant sources (including vegetables such as lettuce, broccoli, wheat, and alfalfa), from the central nervous system of animals, and from a variety of macro- and micro-marine algae. However, monogalactosyldiacylglycerides containing the polyunsaturated fatty acid eicosapentaenoic acid ("EPA") are only found in marine algae. More specifically, monogalactosyldiacylglycerides with the highest content of eicosapentaenoic acid are found in cold water marine micro-algae species. These eicosapentaenoic acid bearing glycerides (i.e., MGDG-EPA) are formed along with many other algal products, thus making the purification and isolation of useful quantities of these materials complex and burdensome.

Accordingly, there is a need in the art for a process for making compositions containing high concentrations of MGDG-EPA. In addition, there is a need in the art for methods relating to the use thereof. The present invention fulfills these needs, and provides further related advantages.



Summary of the Invention

Briefly stated, in one embodiment of the present invention, an anti-inflammatory composition comprising a high content of MGDG-EPA is disclosed. As used herein, "MGDG-EPA" is a monogalactosyldiacylglyceride wherein at least one of the acyl moieties is eicosapentaenoic acid esterified to the glycerol backbone, as represented by the following structure I:




I

1,2-diacyl-(β-D-galactopyranosyl(1'→3))-sn-glycerol

wherein  $R^1$  and  $R^2$  represent a hydrocarbon chain of a fatty acid containing from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds, and wherein at least  $R^1$  or  $R^2$  is the hydrocarbon chain of eicosapentaenoic acid. MGDG-EPA is present in the composition in an amount ranging from 35 to 95 percent by weight of the total composition. In a preferred embodiment, both  $R^1$  and  $R^2$  are the hydrocarbon chain of eicosapentaenoic acid.

In a further embodiment, the anti-inflammatory compositions of the present invention further comprise DGDG. As used herein, "DGDG" is a digalactosyldiacylglycerol as represented by the following structure II:



II

1,2-diacyl-( $\alpha$ -D-galactopyranosyl-(1'→6'))- $\beta$ -D-galactopyranosyl-(1'→3))-sn-glycerol

wherein R<sup>1</sup> and R<sup>2</sup> represent the hydrocarbon chain of a fatty acid containing from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds. In addition, the anti-inflammatory compositions of the present invention may further comprise phosphatidyl choline ("PC"), as well as chlorophylls and carotenoids.

In yet another embodiment, a process of manufacturing an anti-inflammatory composition enriched with MGDG-EPA is disclosed. The process includes the steps of extracting marine algae with an extraction solvent, followed by phase separation with a first solvent protocol to yield an organic phase; fractionating the organic phase by polar chromatographic separation with a second solvent protocol to yield an MGDG-EPA enriched fraction; further fractionating the MGDG-EPA enriched fraction by non-polar chromatographic separation with a third solvent protocol to yield a further enriched MGDG-EPA; and removing solvents from the further enriched MGDG-EPA fraction to yield the anti-inflammatory composition.

In still a further embodiment, a method for treating inflammation comprising topically administering to an animal, including humans, an effective amount of a composition comprising a high content of MGDG-EPA is disclosed. The composition may be administered in various

forms, including emollients, ointments, capsules, tablets, drops, syrup, lozenges, suppositories, inclusions and aerosols.

Other aspects of the present invention will become evident upon reference to the attached figures and following detailed description.

#### Brief Description of the Drawings


Figure 1 illustrates the components of a representative composition of this invention as resolved by thin layer chromatography (TLC).

Figure 2 presents the proton NMR spectra for MGDG-EPA.

Figure 3 compares the anti-inflammatory activity of a representative composition of the present invention (solid line) to that of a control (dashed line).

#### Detailed Description of the Invention


The present invention is directed to an anti-inflammatory composition containing a high content of MGDG-EPA. MGDG-EPA is identified above as structure I, wherein  $R^1$  and  $R^2$  represent a hydrocarbon chain of a fatty acid, and wherein at least  $R^1$  or  $R^2$  is the hydrocarbon chain of eicosapentaenoic acid ("EPA" - an omega-3 fatty acid containing 20 carbon atoms and 5 double bonds). For example, when  $R^1$  of structure I is the hydrocarbon chain of eicosapentaenoic acid and  $R^2$  represents the hydrocarbon chain of a non-EPA fatty acid, the corresponding MGDG-EPA has the following structure III:



III

Similarly, when both R<sup>1</sup> and R<sup>2</sup> of structure I is the hydrocarbon chain of EPA, the corresponding MGDG-EPA has the following structure IV:

5



IV

As used herein, the term "fatty acid" refers to a class of organic compounds containing a saturated or unsaturated, branched or unbranched, substituted or unsubstituted, hydrocarbon chain which terminates with a carboxyl group. The hydrocarbon chains of the fatty acids of the present invention preferably contain from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds, and more preferably from 12 to 22 carbon atoms. Representative examples of fatty acid include (but are not limited to) lauric acid (12:0), myristic acid (14:0), palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1) and eicosapentaenoic acid (20:5).

In a preferred embodiment of MGDG-EPA, both R<sup>1</sup> and R<sup>2</sup> of structure I above are the hydrocarbon chain of EPA. In a further preferred embodiment, the carbon-carbon double bonds of EPA are located at the Δ5,8,11,14, and 17 positions (i.e., *cis*-Δ-5,8,11,14,17-eicosapentaenoic acid).

As indicated above, the compositions of the present invention contain a high content of MGDG-EPA. Specifically, the composition may contain MGDG-EPA in an amount ranging from 35 to 95 percent by weight of the total composition, preferably from 40 to 80 percent by weight of the total composition, and most preferably from

50 to 70 percent by weight of the total composition. In addition to the values specifically identified above, all values falling within the above ranges are expressly incorporated herein.

5           The anti-inflammatory composition of the present invention may further include DGDG as identified above in structure II, wherein  $R^1$  and  $R^2$  represent the hydrocarbon chain of a fatty acid containing from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds. DGDG  
10 may be present in the composition in an amount up to about 20 percent by weight of the total composition.

          The anti-inflammatory composition of the present invention may further include phosphatidylcholine ("PC") in an amount up to 15 percent by weight of the total  
15 composition, as well as minor components of chlorophylls and/or carotenoids.

          Eicosapentaenoic acid ("EPA") is present as a constituent, in an esterified form, of the monogalactosyl-diacylglycerol of the present invention. Although not  
20 intending to be limited to the following theory, the anti-inflammatory function of EPA is believed to involve the chemical messenger system responsible for stimulation of many of the physiological responses to inflammatory challenge, for example, reduced chemotaxis of inflammatory  
25 polymorphonuclear leukocytes. The chemical messenger system is part of the arachidonic acid ("AA") metabolic cascade, often referred to as the eicosanoid cascade. The cascade includes two enzyme systems which act upon AA to produce various prostaglandins, thromboxanes, leukotrienes  
30 and the like, which are important mediators of inflammation. Because of EPA's structural similarity to AA, EPA also is a substrate metabolized by the cascade. A structural difference between the two is the presence of an additional carbon-carbon double bond in EPA. Thus, the  
35 metabolites derived from EPA also retain the additional double bond not present in the metabolites of AA. As a

consequence, EPA metabolites greatly diminish, and in some instances eliminate, the physiological effectiveness of these metabolites relative to their AA counterparts. In addition, EPA is a competitive inhibitor of a key enzyme  
5 in the production of inflammatory prostaglandins and thromboxanes.

A primary step in inflammation is the release of AA from AA-phospholipid reservoirs found in cellular membranes. Exposure to EPA results in the replacement of  
10 AA and a concomitant increase of EPA-bearing phospholipid reservoirs. Upon inflammation challenge, the cascade commences with the release of AA or, if present, EPA by the enzyme phospholipase A<sub>2</sub> which is less reactive with EPA-phospholipids than AA-phospholipids. Inflammation  
15 occurs as a result of AA metabolism whereas the metabolism of EPA does not result in inflammation. Prophylactic EPA therapy to prevent inflammation is based upon the establishment of membrane phospholipid reservoirs enriched with EPA. The composition of the present invention  
20 provides a source of EPA as a substitute substrate in the eicosanoid cascade, thereby inhibiting or preventing inflammation.

The composition may be topically administered either prophylactically or in response to an inflammatory  
25 condition in an amount sufficient to prevent or diminish inflammation. Formulations for topical administration include tablets, capsules, lozenges, suppositories, syrups, drops, ointments, creams, lotions, gels, oils, emollients, inclusions and aerosols. Topical  
30 administration is effective for the prevention or treatment of localized inflammatory conditions. For example, the composition may be administered as an emollient for sunburn, athlete's foot, acne, pruritis, eczema, psoriasis, atopic dermatitis, rheumatoid  
35 arthritis, and skin cancer; as an aerosol for the treatment of bronchial inflammation and esophagitis; as an

enteric coated pill or tablet for inflammatory bowel disease or suppository for rectal inflammation; as a lozenge for esophagitis; and as drops for inflammation of the eye, ear or nose.

5           The composition of the present invention is applied at a concentration and frequency which provides effective prevention or treatment of inflammation. The formulation of the composition, as well as administered concentrations and schedules, can be determined by one  
10 skilled in the art, and will be depend on a number of factors, including the avenue of administration, location and severity of inflammation and composition formulation. In the practice of this invention, the composition containing a high content of MGDG-EPA is preferably  
15 applied at a formulation concentration ranging from 0.1 to 10 percent by weight of the formulation.

          The efficacy of the anti-inflammatory compositions of this invention may be determined by appropriate assays. For example, a preferred general  
20 efficacy assay for topical anti-inflammatory compositions is the mouse ear assay. In the mouse ear assay, the effectiveness of a composition in preventing inflammation is determined by measuring the mouse ear swelling in response to inflammation challenge. In the test, mouse  
25 ears are first treated with the anti-inflammatory composition and then subjected to exposure to a pro-inflammatory composition. The determination of the efficacy of the anti-inflammatory composition of the present invention by the mouse ear assay is described in  
30 Example 3 below.

          The anti-inflammatory compositions of the present invention may be prepared from algae by sequential extraction, polar chromatography, and non-polar chromatography as disclosed in greater detail below. The  
35 genera of algae that may be extracted include (but are not limited to) Chlorella, Chaetoceros, Cyclotella,

Ellipsoidon, Isochrysis, Nannochloris, Nannochloropsis, Nitzschia, Phaeodactylum, Porphyra, Porphyridium, Skeletonema, Thalassiosira, Gigartina, Monochrysis and Monoraphidium. Preferred species within the above genera include Chlorella minutissima, Chaetoceros gracilis, Chaetoceros muelleri, Cyclotella cryptica, Isochrysis galbana, Nannochloropsis salina, Nitzschia dissipata, and Phaeodactylum tricornutum

The marine algae, from which the anti-inflammatory compositions of the present invention are obtained, are first seed cultured in small vessels (such as 15 L carboys) and then pilot scale cultured in shallow tanks, bioreactors, or ponds. The culture medium may be salinated freshwater or preferably sea water. The medium is preferably supplemented with nutrients, including (but not limited to) ammonium, bicarbonate, phosphate, iron, nitrate and trace minerals. Protocols for the large scale propagation of algae are described in, e.g., A. Richmond, Handbook of Microalgal Mass Culture, CRC Press, Boca Raton, FL (1986) (incorporated herein by reference in its entirety). The algae may be harvested once the density of the algal culture is sufficiently high, preferably when the optical density of the algal culture at 690 nm is 1.0. Harvesting may be accomplished by coarse screening of the algae and then concentration of the resulting algal slurry. Concentration of the algae by removal of water is typically accomplished by centrifugation, sedimentation, evaporation, flocculation, ultrafiltration, flotation, or a combination of these techniques. The resulting dewatered algae are suitable for extraction.

The algae are then extracted with an extraction solvent(s) to yield an extraction solvent extract. The extraction solvent or solvents solubilize the MGDG-EPA component(s) present in the algae. Suitable extraction solvents include polar organic solvents such as alcohols, esters, ethers, ketones, and aldehydes, mixtures thereof,



and mixtures thereof containing water. Preferred extraction solvents include methanol, ethanol, propanol, isopropanol, acetone. In a preferred embodiment, the extraction solvent is 90% (i.e., 180 proof) aqueous ethanol. The algae may be extracted with an extraction solvent at ambient temperature or preferably near the boiling point of the extraction solvent. The extraction process includes contacting the algae with the extraction solvent for a period of time sufficient to effect solvent solubilization of the MGDG-EPA components present in the algae. For example, extraction with 90% (180 proof) aqueous ethanol at a temperature near its boiling point for 10 minutes is preferred. Upon cooling the mixture, the ethanolic extract containing MGDG-EPA is separated from the residual algal solids by filtration to yield the extraction solvent extract.

The extraction solvent extract is then fractionated by phase separation with a first solvent protocol comprising an organic water-immiscible solvent and water. The organic water-immiscible solvent selectively solubilizes the MGDG-EPA present in the extraction solvent extract while the water serves to remove highly polar components from the extraction solvent extract such as sugars, amino acids, and other water soluble components. Suitable organic water-immiscible solvents include non-polar organic solvents such as hydrocarbons, ethers, and chlorinated hydrocarbons among others. Preferred organic water-immiscible solvents include pentane, hexane, heptane, petroleum ethers, diethyl ether, dichloromethane, and chloroform among others. In a preferred embodiment, the organic water-immiscible solvent is hexane. After thorough agitation of the extraction solvent with the organic water-immiscible solvent and water, the aqueous phase is separated from the organic phase. Removal of the organic solvents from the organic phase yields a crude algal lipid extract

containing the MGDG-EPA components. In a preferred embodiment, the extraction solvent and first solvent protocol comprises 90% aqueous ethanol, hexane, and water in a ratio of 1:1:1.

5           The crude algal lipid extract containing MGDG-EPA is then fractionated by polar chromatography. A solution of the crude algal lipid extract in a suitable solvent, such as 10% ethanol in hexane, is applied to a polar chromatographic column. The solid phase of the  
10 polar chromatographic column may be alumina, deactivated alumina, silicic acid, Florisil®, DEAE cellulose, or other suitable polar solid phase. In a preferred embodiment, the polar solid phase is deactivated alumina (described in greater detail in Example 1 below). The crude algal lipid  
15 extract is fractionated by polar chromatography by elution with a second solvent protocol. The second solvent protocol may be a suitable fractionating solvent or mixtures of solvents including water and mixtures of polar and non-polar organic solvents in various proportions.  
20 Suitable polar organic solvents include alcohols, esters, ethers, ketones, and aldehydes among others. Preferred polar organic solvents include alcohols and esters such as methanol, ethanol, and ethyl acetate. Suitable non-polar solvents include hydrocarbons, petroleum ethers, and  
25 chlorinated hydrocarbons among others. Preferred non-polar organic solvents include pentane, hexane, diethyl ether, dichloromethane, and chloroform. In preferred embodiment, the polar organic solvent is ethanol and the non-polar organic solvent is hexane. In a preferred  
30 embodiment, the second solvent protocol comprises water, ethanol, and hexane. In a typical polar chromatography, the column is eluted with increasingly polar solvents or solvent mixtures. For example, a polar column may be sequentially eluted with 10% ethanol in hexane, 40%  
35 ethanol in hexane, 100% ethanol, and 20% water in ethanol. When the crude algal lipid extract is fractionated by

deactivated alumina as the solid phase and with the above elution pattern, the fraction eluted with 20% water in ethanol contains the MGDG-EPA enriched fraction.

The enriched MGDG-EPA fraction is then further  
5 fractionated by non-polar chromatography to yield a further enriched MGDG-EPA fraction. Specifically, the enriched fraction, in a suitable solvent such as ethanol/water (1:1), is applied to a non-polar chromatographic column. The solid phase of the  
10 chromatographic column may be any suitable non-polar or reverse phase solid phase. In a preferred embodiment, the non-polar solid phase is reverse phase silica, and in a most preferred embodiment, the non-polar solid phase is reverse phase octadecylsilyl silica (ODS). The further  
15 enrichment of the MGDG-EPA fraction is accomplished by elution with a third solvent protocol. The third solvent protocol may be any suitable fractionating solvent or mixtures of solvents including water and mixtures of polar organic solvents in various proportions. Suitable polar  
20 organic solvents include alcohols, esters, ethers, ketones, and aldehydes among others. Preferred polar organic solvents include alcohols and esters such as methanol, ethanol, and ethyl acetate. In a preferred embodiment, the polar organic solvent is ethanol and the  
25 third solvent protocol comprises water and ethanol in varying proportions. In a typical non-polar chromatography, the column is eluted with decreasingly polar solvents or solvent mixtures. For example, a non-polar column may be sequentially eluted with 50% (100  
30 proof) aqueous ethanol, 70% (140 proof) aqueous ethanol, 80% (160 proof) aqueous ethanol, 90% (180 proof) aqueous ethanol, and 100% (200 proof) ethanol. When the enriched MGDG-EPA fraction obtained from polar chromatographic separation was further fractionated by non-polar  
35 chromatography with ODS as the solid phase and with the

above solvent system, the further enriched MGDG-EPA fraction eluted with 90% (180 proof) aqueous ethanol.

The following examples are offered by way of illustration, not limitation.

5

### EXAMPLES

#### Example 1

This example illustrates the preparation of an  
10 anti-inflammatory composition of the present invention.

The marine microalga Chlorella minutissima was grown in cultivation tanks 4' x 10' with a paddle wheel system for agitation and a culture medium depth of 10"-12". The culture medium consisted of seawater,  
15 supplemented with ammonium nitrate (8 mM), sodium bicarbonate (5 mM), potassium dihydrogen phosphate (0.04 mM), and iron versenate (0.02 mM). The cultures were grown in full sunlight, or in plastic covered green houses, and the nutrient levels were measured weekly and  
20 additional nutrients added as required. When the cultures reached an optical density of 1.0 (measured at 690 nm) the algae were harvested. Fifty percent of each of 6 culture tanks was harvested giving a total of approximately 700 gallons.

25 The culture was pumped through a mixing tower where metered amounts of coagulant (Calgon WT-2511) were added with rapid agitation. The mixture next passed into a non-stirred column where metered amounts of flocculent (Betz Polymer 1160) were added, and thence into settling  
30 tanks. Settling was rapid and a large proportion of supernatant water was discarded. After further settling overnight, in excess of about 90% of the supernatant liquid was discarded, leaving a total volume of approximately 25 gallons. To this was added 3.64 kg of  
35 diatomaceous earth (DE) as a filter aid. Further rapid settling occurred, thus allowing the removal of more

supernatant water to leave 12 gallons of slurry. The slurry was vacuum filtered to yield a cake weighing 11.4 kg. The dry weight of algae contained in the cake was 686 gm.

5           The algal cake was extracted in 1.5 kg aliquots by adding to 4 L of boiling 90 percent aqueous ethanol, bringing the mixture back to a boil (5-10 minutes), and then continuing boiling for 10 minutes. The extract was obtained by filtration through a Buchner funnel under  
10 vacuum to remove solids. Water and hexane were added to the cooled extract to provide a final proportion (extract/hexane/water) of 1:1:1 and the mixture was agitated well and allowed to separate into two phases. The lower, aqueous ethanol phase contained water soluble  
15 materials, including sugars, amino acids, and some pigments. The upper hexane phase contained the crude lipid extract. The solvent was evaporated from the upper phase and the residue taken up in a minimal amount of 10% ethanol in hexane. At this stage small aliquots were  
20 taken for determination of dry weight, for thin-layer chromatography, and for fatty acid analysis by transesterification and gas chromatography. Total lipid extract was 98.8 gm, or 14.4% of dry weight of algae.

          The crude lipid extract, dissolved in 10%  
25 ethanol in hexane to a total volume of 880 ml, was applied to a 4" x 24" alumina column prepared as follows: activated alumina (aluminum oxide) of a chromatographic grade (Aldrich Chemical Co., Brockman grade I, acidic, about 150 mesh) was deactivated by suspending it overnight  
30 in two volumes aqueous ethanol at a mixture of 70 parts ethyl alcohol to 30 parts water. The deactivated alumina was then packed into the column as a slurry in the water/alcohol mixture. The column was washed well with four volumes of absolute ethanol to remove the water,  
35 followed by a wash with one column volume of 40% ethanol in hexane and two column volumes of 10% ethanol in hexane.

The crude lipid extract was pumped onto the column and eluted with 4 L of 10% ethanol in hexane, 3 L of 40% ethanol in hexane, 4 L of 100% ethanol, 4 L of 20% water/80% ethanol and 3 L ethanol wash. All fractions  
5 were vacuum evaporated, the residues taken up in 10% ethanol in hexane and samples applied to TLC plates for analysis of lipid composition.

The 20% water/80% ethanol fraction, which contained the MGDG-EPA, was evaporated to dryness (weight  
10 - 12.4 gm) and taken up in ethanol/water 1:1. The MGDG enriched fraction was then further separated on a 2" x 12" reverse-phase column of ODS (Preparative C18, 125A, Millipore Corp.) which was prepared as follows: the ODS-silica was slurried with absolute ethanol and packed in  
15 the column in ethanol. After a wash with 1 L of absolute ethanol, the column was washed with 1 L of 75% ethanol/25% water, then 1.5 L of ethanol/water 1:1. The MGDG-EPA-containing fraction from the alumina column was divided in two, and one portion (~6.2 gm) at a time  
20 separated on the ODS column in two separate runs.

Elution was with 1 L of 50% aqueous ethanol, 1 L of 70% (140 proof) aqueous ethanol, 1.5 L of 80% (160 proof) aqueous ethanol, 1 L of 85% (170 proof) aqueous ethanol, 1.5 L of 90% (180 proof) aqueous ethanol and 2.5  
25 L of absolute ethanol. In each run, the 90% aqueous ethanol fraction contained the bulk of the MGDG-EPA (approx. 90% of the total) as determined by TLC. These 90% fractions from the two runs were combined and evaporated (total weight 10.2 gm). The composition of the  
30 combined fractions was approximately 60% MGDG-EPA, 30% eicosapentaenoic acid believed to be bearing sulphoquinovosyldiacylglyceride ("SQDG-EPA"), 7% DGDG and 3% PC (with minor components of carotenoid and chlorophyll).

Example 2

This example illustrates the characterization of the MGDG-EPA components of Example 1.

A sample of the composition containing MGDG-EPA of Example 1 was subjected to preparative thin layer chromatography separation. Two major components were observed, MGDG-EPA and SQDG-EPA, as identified in Figure 1. MGDG-EPA was removed from the plate by scraping followed by elution from the silica. The proton NMR spectra of the MGDG-EPA component is presented in Figure 2. In addition, a portion of the MGDG-EPA was transesterified and the resulting fatty acid esters were analyzed by gas chromatography. The identity of the components of the fatty acid ester mixtures was made by comparison to retention time with known standards run immediately before and after the samples, and by carbon number plotting. The fatty acids of the MGDG-EPA component are (in percent by weight) presented in Table 1 below:

Table 1

|    | <u>Fatty</u> | <u>Percent</u> |
|----|--------------|----------------|
|    | <u>Acid</u>  | <u>MGDG</u>    |
| 25 | 12:0         | tr             |
|    | 14:0         | 5.4            |
|    | 16:0         | 0.2            |
|    | 16:1         | 1.4            |
|    | 16:5         | ---            |
| 30 | 18:1         | ---            |
|    | 20:2         | ---            |
|    | 20:4         | 1.5            |
|    | 20:5         | 91.5           |
|    | 22:4         | ---            |

In Table 1 above, the first number of the fatty acid represents the number of carbon atoms in the fatty acid, and the number following the colon represents the number of double bonds in the hydrocarbon chain of the fatty acid. The identity of the major fatty acid constituent (i.e., 20:5) as eicosapentaenoic acid was confirmed by mass spectrometry. In the practice of this invention, EPA constitutes 50-99% of the fatty acids esterified to MGDG, preferably from 70-95%, and more preferably from 80-90%.

### Example 3

This example illustrates the efficacy of a composition of the present invention as an anti-inflammatory agent in the generic mouse ear test.

Both ears of mice were treated with either petrolatum (Vaseline®) or a formulation of petrolatum and a representative composition of this invention containing 85% by weight MGDG-EPA, 15% by weight DGDG, and trace chlorophylls (with EPA totaling 50% by weight of the total composition). The formulation used in this study contained 2% by weight of the above composition and 98% by weight petrolatum. Treatment involved the application of 5 mg of either petrolatum by itself or the above 2% formulation. Application was accomplished with a wooden stick to each ear at 9:00 a.m. and 5:00 p.m. for 5 consecutive days. Five mice (i.e., 10 ears) were used for both the control (i.e., petrolatum only treatment) and test animals (i.e., animals treated with the above 2% formulation). Two hours after the last application, 10  $\mu$ l of 10% croton oil in acetone was applied to each ear to induce inflammation. The total thickness of the ears was measured with a micrometer having a measuring surface of 1 cm diameter, which is approximately the size of the ear. Measurements were taken immediately before challenge (t = 0) and 2, 4, 8, and 24 hours after croton oil




application. The results of this experiment are presented in Figure 3, with ordinate representing the change in ear thickness in microns (i.e., swelling) as determined from  $t = 0$  to  $t = 2, 4, 8$  and 24 hours. Figure 3 illustrates a statistically significant depression of ear swelling by the 2% formulation (solid line) at 4, 8, and 24 hours after induction of inflammation compared to the petrolatum control (dashed line).

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except by the appended claims.

Claims

1. An anti-inflammatory composition comprising a monogalactosyldiacylglycerol having the structure:




wherein  $\text{R}^1$  and  $\text{R}^2$  represent a hydrocarbon chain of a fatty acid containing from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds, wherein at least  $\text{R}^1$  or  $\text{R}^2$  is the hydrocarbon chain of eicosapentaenoic acid, and wherein the monogalactosyl-diacylglycerol is present in the composition in an amount ranging from 35 to 95 percent by weight of the total composition.

2. The composition of claim 1 wherein both  $\text{R}^1$  and  $\text{R}^2$  are the hydrocarbon chain of eicosapentaenoic acid.

3. The composition of claim 1 wherein the monogalactosyldiacylglycerol is present in the composition in an amount ranging from 40 to 80 percent by weight of the total composition.

4. The composition of claim 1 wherein the monogalactosyldiacylglycerol is present in the composition in an amount ranging from 50 to 70 percent by weight of the total composition.

5. The composition of any one of claims 1-4 further comprising a digalactosyldiacylglycerol having the structure:



wherein  $R^1$  and  $R^2$  represent the hydrocarbon chain of a fatty acid containing from 10 to 25 carbon atoms and from 0 to 6 carbon-carbon double bonds.

6. The composition of any one of claims 1-5 further comprising phosphatidylcholine.

7. A process of manufacturing an anti-inflammatory composition an MGDG-EPA, comprising:

extracting algae with an extraction solvent to yield an extract;

phase separating the extract with a first solvent protocol to yield an organic phase;

fractionating the organic phase by polar chromatographic separation employing deactivated alumina with a second solvent protocol to yield an enriched diglyceride fraction;

fractionating the enriched diglyceride fraction by non-polar chromatographic separation with a third solvent protocol to yield a further enriched diglyceride fraction; and

removing solvents from the further enriched diglyceride fraction to yield the anti-inflammatory composition.

8. The process of claim 7 wherein the algae is selected from the genera Chlorella, Chaetoceros, Cyclotella,

Ellipsoidon, Isochrysis, Nannochloris, Nannochloropsis,  
Nitzschia, Phaeodactylum, Porphyra, Porphyridium,  
Skeletonema, Thalassiosira, Gigartina, Monochrysis and  
Monoraphidium.

9. The process of claim 7 wherein the alga is Chlorella minutissima.

10. An anti-inflammatory composition made according to the process of claim 7.

11. A method for treating inflammatory disorders comprising topically administering to an animal an effective amount of a composition comprising MGDG-EPA, wherein MGDG-EPA is present in the composition in an amount ranging from 35 to 90 percent by weight of the total composition.

12. The method of claim 11 wherein the composition is administered in the form of an emollient, syrup, oil, tablet, capsule, lozenge, aerosol, drop, suppository, ointment, or inclusion.




FIG. 1

MGDG-EPA

Fig. 2




FIG. 3.

